



Orius 3.6F Foliar Fungicide
For Use on Barley to suppress *Fusarium* head blight (scab)

SECTION 18 SPECIFIC EXEMPTION
EE#: 07-MT-05

**FOR DISTRIBUTION AND USE ONLY IN THE STATE OF
MONTANA
UNDER SECTION 18 SPECIFIC EXEMPTION**

These use directions must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire Directions for Use and Disclaimer of Warranties before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for Orius 3.6F Foliar Fungicide (EPA Registration Number 264-752-66222).

Any adverse effects resulting from the use of Orius 3.6F Foliar Fungicide under this specific exemption must be immediately reported to the Montana Department of Agriculture.

EFFECTIVE PERIOD:

These use directions become effective May 15, 2007 and expire on July 20, 2007.

DIRECTIONS FOR USE

A maximum of 45,500 acres of barley may be treated under this exemption.

RECOMMENDED APPLICATIONS

CROP	DISEASES	RATE OF ORIUS 3.6F
Barley	<i>Fusarium</i> head blight (scab)	4 fl oz per acre
Apply Orius 3.6F as a preventive foliar spray when the earliest disease symptoms appear on the leaves or stems. For optimum disease control, the lowest rate of a spray surfactant should be tank-mixed with Orius 3.6F. Orius 3.6F may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 3 gallons of spray solution per acre by aircraft spray equipment.		
Application through any type of irrigation equipment is prohibited.		
A maximum of 1 application per season may be made. A maximum of 4 fl oz of Orius 3.6F may be applied per acre per crop season. Orius 3.6F may be applied up through the beginning of flowering (Feekes 10.5). Application may not be made within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Do not enter treated field for 12 hours after application.		
Do not make more than 1 application of tebuconazole-containing product to barley per season.		

ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater, estuarine, and marine fish and invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Do not apply by ground or air within 100 feet of aquatic areas such as lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds and estuaries.

For transportation emergencies, call INFOTRAC 24 hours a day: 1-800-535-5053

For medical emergencies, call PROSAR 24 hours a day: 1-877-250-9291

For product information, call MANA: 1-866-626-2462

Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Rd., Suite 300
Raleigh, NC 27609